


Other Information

Bone Marrow Registry

The **IBTS** also runs the Irish Unrelated Bone Marrow Registry which facilitates stem cell transplants for Irish and International patients. If you are interested in joining the bone marrow registry please ask for an information leaflet on clinic, be sure to indicate your interest on the back of the HLQ and inform a staff member before the donation process is started. Further information on bone marrow donation is also available at **giveblood.ie/bonemarrow_marrow_registry**.



Use of Blood Components and Samples for Research and Development

Donated blood is separated into red cells, platelets, plasma and occasionally granulocytes (white cells) which are blood components. The vast majority of red cells and platelets is transfused directly to patients e.g. after blood loss or cancer treatment.

The **IBTS** on occasions uses parts of donated blood for laboratory work, education, training or research purposes, within the **IBTS**, the Health Service or the Universities/Institutes of Technology. This includes white cells which are not given to patients, or blood not used within its shelf life. The **IBTS** does not receive any payment for this. Every time donated material is used for these purposes, the donation identification number, blood group, and occasionally the donor's age and gender, will accompany the donation. This personal data is required for traceability purposes and research analysis. Please visit **https://healthprofessionals.giveblood.ie/research/donated-material-for-health-research/** to learn more about how your donation supports health research. In order to deliver a high standard quality service, the **IBTS** also, when required, uses blood within its shelf life to validate/quality check new blood bags or to test kits and equipment.

The **IBTS** also collaborates with some commercial entities to provide remnants of **IBTS** donor samples for research and development studies such as evaluating performance of manufacturers' assays, evaluating materials and qualification of product changes. The **IBTS** and all blood services rely on commercial tests to ensure safe blood and collaborate in improving assay performance characteristics with commercial companies.

If parts of your samples or donations are used for the above purposes, we will ensure, where appropriate, that there is no direct link between your personally identifiable data and the donated material and that approval from a research ethics committee is obtained where necessary.

The **IBTS** considers that this use of donations/samples is appropriate. The **IBTS** receives payment from some of these collaborations which helps offset some **IBTS** costs and is therefore of benefit to the Irish Health Service. The testing and use of your blood in the circumstances outlined above is of benefit to patients and by signing the Donor Declaration on the questionnaire you are giving the **IBTS** consent to do so where required.

We would also like our donors to know that at times their donations are not used. This could happen if the donation was not a complete donation or if the donation is not used before its expiry date.

BT - 0353 [9]

Data Protections and use of personal data

The **IBTS** is committed to protecting the confidentiality of donors and to meeting our responsibilities under the Data Protection Acts 1988 – 2018 and the General Data Protection Regulation 679/2016 (GDPR).

We process your information in order to assess your eligibility to donate; to communicate with you; to ensure both donor and patient safety; for clinical and regulatory audit purposes; to provide safe and suitable blood and tissue products to patients and to ensure the traceability of all donations in line with our legal requirements.

We also process anonymised data for research purposes.

Your information is managed appropriately and securely at all times, and we are required to retain records for 30 years.

For more information on the **IBTS** Data Protection Policy and your Data Protection rights please ask for our leaflet on clinic or see **www.giveblood.ie/privacy**

Communication

As part of the donation process you will be asked to consent to the **IBTS** contacting you after donating and ahead of future donations.

Texting is the primary method of communication we use pre and post clinic.


Phone calls are used to answer donor queries; provide medical follow-up where needed and for specific requests to attend clinics;

Letters are also used to provide medical follow-up when required.

Email is used to answer donor queries; communicate about service evaluation and research projects; invite donors to participate in surveys and keep donors up to date about new developments.

Social media platforms are used to communicate with donors, sharing general information on blood donation and inspirational stories about donors and patients.

For more information on the **IBTS** communication policy please ask for our leaflet in clinic or log onto **www.giveblood.ie** and search Privacy. If you have any further questions please ask one of our staff on the clinic.



THANK YOU

for attending this **Blood Donation** Clinic today

BLOOD SAFETY – KEEP TRANSFUSIONS SAFE

Help us keep the blood supply safe for patients by reading this information before you donate.

All donations are tested for infections that can be passed on by blood or platelet transfusions. **Early stage infection may not always show up on testing.** This is why we must take great care in donor selection and why you must not donate to see if you are infected. If you donate to see if you are infected you are putting patients' lives at risk.

We rely totally on our donors to answer the questions we ask honestly and accurately to the best of their knowledge, as this is essential for the safety of the patients who receive blood. We ask that you please do so. All information you provide is confidential.

You must **NEVER** donate if:

?

You think you need a test for HIV or hepatitis. (You can get a test from your GP or a sexual health clinic)

HIV or Hep

You think your partner needs a test for HIV or hepatitis

HIV

You or your partner has HIV or is receiving treatment for HIV

HTLV

You or your partner has HTLV

Hep B or Hep C


You, your partner or close household contacts have hepatitis B or C*

STI

You have **ever** had or been treated for Syphilis

€

You have **ever** received money or drugs for sex



You have **ever** injected, or have been injected with non-prescribed drugs; **even once or a long time ago.** This includes body building drugs, injectable tanning agents and injected chemsex drugs.

You must **NOT** donate for at least **4 months** after sex if:

?

You have had anal sex with a new sexual partner**, or you had sex with more than one sexual partner and had anal sex

**A new sexual partner is someone you have not had sex with before or a person with whom you resumed a previous sexual relationship in the past 4 months

You must **NOT** donate for at least **4 months** after:

You snorted cocaine or any other drug

Conclusion of treatment and full recovery from Gonorrhoea

You have taken medication to prevent HIV infection (pre /post exposure prophylaxis, PrEP/ PEP)

You have taken part in Chemsex i.e. have used drugs (other than cannabis, alcohol, Viagra or other medication for erectile dysfunction) before or during sex to enhance sexual experience. Chemsex is a type of sexual activity where people take certain drugs to enhance their sexual experiences, and let them have sex for longer or with more people. Drugs associated with chemsex include methamphetamine, mephedrone, GHB and other drugs.

NB: IF you were injected or injected yourself with drugs during Chemsex even once or a long time ago you will never be able to donate

More information about Individual Donor Risk Assessment (IDRA) is available in the IDRA leaflet, please ask a staff member.


Irish Blood Transfusion Service

Seirbhís Fulaistriúcháin na hÉireann

BT - 0378 [9]

Donation Information for Platelet Donors

Please read this donation information carefully, you will be asked to sign a donor declaration to confirm that you have read and understood this information and consent to the donation process.



giveblood.ie

*There are exceptions so please speak to a nurse.

Registration (New Donor)

In order to create a unique identification for you in the **IBTS** database, you will be asked to provide photo ID and give your name, date of birth, contact details, phone number and email address. This information will be used to contact you and send reminders about clinics. You will complete a Health and Lifestyle Questionnaire (HLQ). This asks specific questions about your health, lifestyle and travel history.

Interview – Health Screening (New Donor)

As a new donor or a donor who has not donated in the last 5 years, you will be interviewed by a Nurse on the clinic following completion of the HLQ that you were given at registration. This is to make sure that you are well enough to donate and that your donation is safe to give to a patient. Information for new donors born outside of Ireland / the United Kingdom will be provided in a separate leaflet.

On some occasions it may not be possible to accept a donation from you for either medical or other reasons, this will be explained, and you will be told if and when you are eligible to attend a clinic again.

As an apheresis donor, only samples are taken following interview on your first ever attendance or if your last donation was greater than 5 years ago. You will also be given additional information by the nurse during the interview to read.

Registration (Regular Donor)

As you have previously registered with the **IBTS** you will be asked to provide photo ID and to confirm your personal details, then you will be asked to complete a Health and Lifestyle questionnaire (HLQ).

Interview – Health Screening (Regular Donor)

On completion of the HLQ you will have an interview with a staff member to assess your eligibility to donate. On some occasions it may not be possible to accept a donation from you for either medical or other reasons, this will be explained and you will be told if and when you are eligible to attend a clinic again.

Consent

You will be asked to read and sign a Donor Declaration on the HLQ following your interview. Please read this carefully as this provides the IBTS with permission to:

- Take a donation
- Carry out testing (HIV, Hepatitis and other infectious agents outlined below)
- Issue your donation to patients in hospital or for use indirectly as outlined below
- Contact your Doctor or hospital to obtain further information regarding your eligibility to donate, if required.

If you decide at any time that you do not want to go ahead with the donation process you can leave the clinic, please return your questionnaire to a staff member.

Once you have provided your information to the **IBTS** in order to donate, it must be retained so that we can meet our statutory requirements. Please ask our staff if you would like more information on this.

Donation Process

Prior to your first donation we will check your Hb and platelet counts. If on the last platelet donation your counts were noted to be lower than the required levels, we will check them prior to the donation starting. Your blood pressure and pulse will be checked before you donate. A needle will be inserted into a suitable vein in your arm. This needle is attached to a set loaded on to the apheresis platelet collection machine, a cell separator. This set is only used once and disposed of when the donation finishes.

Samples of blood will be taken at the start of the donation from a small pouch attached to the apheresis set. These samples are used for routine testing. We will also test your Hb and platelet levels (if not already done). During the donation, the platelets are separated from the remainder of the blood and periodically your red cells and plasma are returned back to you. The process takes an average of 60 minutes but never more than 100 minutes. You will be asked to apply firm pressure to your arm after the needle has been removed for at least 3 minutes to help prevent bruising.

During the donation if you feel in any way unwell or uncomfortable, please inform a staff member immediately. Staff on clinic observe donors throughout the process and are trained to manage the complications associated with donation.

Donating platelets can lead to iron deficiency. This may be associated with tiredness, impaired concentration or poor work performance, in some people. Continued loss of iron can cause anaemia.

After the Donation

You will be asked to rest for a short while in our canteen and offered refreshments and a snack. Enjoy, you deserve it! You will be given a post donation advice card by staff explaining the dos and don'ts following donation. It is important to follow this advice and remember to contact the **IBTS** as soon as you can if you become unwell in the 4 weeks following your donation. It is also important to contact the **IBTS** as soon as you can if you remember something that could impact a patient in hospital who would receive your platelets as outlined at the interview.

Testing your Donation

Your blood group is checked every time you donate. The **IBTS** also tests every donation for HIV, hepatitis B, hepatitis C, hepatitis E and syphilis. The first time you donate, your blood will also be tested for HTLV (a virus carried in blood that can cause serious blood and nervous system diseases). The **IBTS** may also carry out extra tests on some donations, for e.g. if you have travelled to a West Nile Virus at risk area, or to meet the needs of certain patients such as babies.

If a positive result for any of these blood tests is found, the donation will be discarded and not used for a patient. You will be contacted by the **IBTS** and offered advice and further testing.

The **IBTS** is legally bound to inform the Public Health Service if your tests confirm that you are positive for **HIV, hepatitis B, C, E or syphilis**. Your personal details, name, address and the results of the blood tests will be sent to the Public Health Service. In a small number of donors, the blood results can give a false positive reaction. You will be informed if this occurs and while there are no implications for your health, we will not be able to use the donation for patients. We freeze a small sample of your blood and retain it for 10 years in case it is necessary to carry out additional tests at a later date.

A+

A-

B+

B-

AB+

AB-

O+

O-

Possible Complications of Donation

For the majority of donors the process is a simple and trouble free experience, however in some cases problems may occur during or after the donation. If you have any problems after leaving the clinic please contact the **IBTS** on 01 4322800 (Dublin) or 021 4807400 (Cork) or contact your GP. **All rates quoted are from IBTS data 2017*

Bruising

Sometimes bruising can occur either during or after the donation (1/279). If noted during the donation, it will be stopped and a staff member will give you an information leaflet with advice about how to best manage the bruising e.g. avoid heavy lifting or manual work where possible for at least 12 hours. If you notice bruising after a donation, report this to a staff member, or if at home apply a cold pack or ice in a bag with a layer of protection over the skin by using a cloth and again avoid heavy lifting or manual work as outlined above. You may experience discomfort associated with the bruising.

Nerve Injury / Nerve irritation / Pain

You must tell us immediately if you have any of the following when the needle is inserted, during the donation or after the needle is removed (1/813):

- pain
- discomfort
- altered sensation in your arm, hand or fingers

Very infrequently, insertion or removal of the needle may cause irritation (1/3,703) or injury of a nerve in your arm (1/8,333). Pain, discomfort or altered sensation in your arm, hand or fingers may indicate that a nerve has been injured.

These symptoms may also occur if a collection of blood under the skin (bruise) is pressing on a nerve. Tell us immediately if you have any of these symptoms and we will stop the donation and give you the appropriate advice.

Symptoms may occur after you have left the donor clinic. One in 141,429 continues to have pain for more than a year after donating.

Tendon Injury / Pain

On rare occasions, a needle may come into contact with a tendon and cause a sharp pain when the needle is inserted. Tell us immediately if you experience any pain or discomfort when the needle is inserted, and we will stop the donation. No tendon injuries were recorded in the **IBTS** in 2017.

Bleeding from the Needle site

Sometimes blood will start to leak from the needle site after the donation is finished (1/352). If this happens, lift your arm above your shoulder and apply firm pressure until the bleeding stops, a staff member will apply another plaster as required. Leave the plaster on the site until the next day to prevent infection.

Arterial Puncture

Rarely (1/10,000), the needle may inadvertently be inserted into an artery. If this happens the needle will be removed immediately and you will be given immediate treatment and appropriate advice.

BT - 0378 [9]

Fainting / feeling lightheaded

Some donors may feel faint (1/99) after donating and a small number faint (lose consciousness) (1/833). There are a number of things that you and our staff can do to minimise this risk so please follow their advice on clinic. If you were to faint on clinic our staff are trained to look after you and will not let you leave the clinic until you are fully recovered. They will also give you an information leaflet with advice prior to going home. If you were to experience a head injury as a result of a faint you would be referred to hospital from the clinic. If you feel faint after leaving the clinic, lie down straight away and drink plenty of non-alcoholic fluids.

Uncommon risks of donation

We collect around 130,000 donations of blood every year and 20,000 doses of platelets. About once or twice a year a donor may need to be admitted to hospital as a result of a complication of giving blood or platelets. This could occur if for example a donor faints and bangs his / her head resulting in an injury. There is a very remote risk of Stroke or Death; however no one has ever died in Ireland due to blood or platelet donation.

In the unlikely event that you experience a fast / irregular pulse or tightness in your chest, this may not necessarily be caused by the donation. If it does occur during or after donation let us know immediately.

Complications specific to Apheresis

Low lymphocyte counts
A recent study has shown that some regular platelet donors have low lymphocyte counts (lymphopaenia). This may be because these white cells are removed during the apheresis procedure. This has not been associated with any illness in donors.

Haemolysis and Air Embolus
Haemolysis occurs when red cells get broken down in the machine by excessive centrifugal speed, or if they are forced through a narrowing in the apheresis set. An air embolus is the inadvertent entry of air into your circulation. These complications are very rare events, due to the safety measures incorporated in the apheresis technology. They have never occurred in the **IBTS** with the Trima Accel machines that have been in use for the last 20 years.

Generalised Allergic Reaction
Very rarely a platelet donor may experience a generalised allergic reaction to the gas used to sterilise the apheresis set. One platelet donor had a severe acute allergic reaction while donating in the **IBTS** in the last 15 years.

Reaction to the Anticoagulant

An anticoagulant (citrate) is used to prevent the blood from clotting in the disposable set. Apheresis procedures cannot be carried out without it. Many donors experience a tingling sensation around the mouth, a metallic taste or a chilly sensation. These are usually very mild symptoms. You must inform our staff if you feel any of these symptoms. Our staff will be able to intervene in the procedure, usually without stopping it, by slightly slowing the flow of the anticoagulant and the symptoms usually improve. If they do not improve or you find them distressing, tell us and we will stop the procedure (1/294).

Citrate Toxicity

Rarely a donor may experience dizziness, light-headedness, muscle cramps, body vibrations, nausea or vomiting due to the anticoagulant that is used in platelet donations.

Tell our staff immediately if you experience any of these symptoms.

In the interests of your health, the donation will be stopped. On extremely rare occasions citrate may disturb the rhythm of the heart. It is important that these symptoms are detected early to prevent a cardiac complication (1 or 2 platelet donors are taken by ambulance to Accident and Emergency ever year).

BT - 0378 [9]