

Document Detail

	PMF IBTS SPEC IBTS/PMF/SPEC/0214[7]
Title:	PLATELETS, ADULT DOSE WITH PLASMA / PAS,
	IRRADIATED
Owner:	QA DOC CON QA DOC CONTROL
Status	CURRENT
Effective Date:	25-Aug-2021
Expiration Date:	25-Aug-2023

<u>Review</u>

Review: IBTS PMF REVIEW

Level	Owner Role	Actor	<u>Sign-off By</u>
1	DOCUMENT CONTROLLER	DEBBIE MAC RORY	DEBBIE MAC RORY
2	QUALITY CONTROL WRITER NBC	ZUZANNA MICHALSKA	ZUZANNA MICHALSKA
2	QUALITY CONTROL REVIEWER NBC	ENDA CADDEN	ENDA CADDEN
2	QUALITY CONTROL REVIEWER NBC	ELIN JOHANSSON	SMARTCOMM
3	QUALITY CONTROL HEAD OF DEPT NBC	ENDA CADDEN	ENDA CADDEN
3	NATIONAL MEDICAL DIRECTOR	STEPHEN FIELD	STEPHEN FIELD
3	LABS HEAD OF MANUFACTURING & ISSUE IBTS	BARRY DOYLE	BARRY DOYLE
4	QUALITY ASSURANCE REVIEWER IBTS	COLIN JOHNS	COLIN JOHNS

Change Orders

Changes as described on Change Order: <u>Change Order No.</u>

Change Orders - Incorporated

Changes as described on Change Order:

Change Order No. IBTS/CO/0248/21

IRISH BLOOD TRANSFUSION SERVICE

PRODUCT MASTER FILE

AS, Marshall PLATELETS, ADULT DOSE WITH PLASMA / PAS, **TITLE:**

Change Description:

Change lower volume specification from 251ml to 271ml

Reason for Change: As per CC107/21/IBTS

Change order No.: IBTS/CO/0248/21

Referenced Documents N/A

SmartSolve Roles N/A

Training Type N/A

SmartSolve Document Category

Category	Mobile	Cryobiology	Website	GDP
Yes / No	No	No	Yes	No
101				
7				

IRISH BLOOD TRANSFUSION SERVICE PRODUCT MASTER FILE

Platelets, Adult Dose with Plasma / PAS, Irradiated

Name of Product:

Title:

PLATELETS, Adult Dose with Plasma / PAS, Irradiated •

E Progesa Codabar Component Code:

E Progesa ISBT – 128 Component Code:

E6953V00

69740

General Description: Platelets prepared from a pool of four buffy coats suspended in approximately 70% additive solution and 30% plasma. The removal of the majority of leucocytes is achieved by filtration.

General Specification:		
Parameter 58	Quality Requirements	Frequency of Control
Volume	271 to 378 ml	100%
Platelet Content	> 200 x 10 ⁹ /unit	1%
Platelet Concentration	\geq 35 ml per 60 x 10 ⁹ of platelets	1%
Leucocyte Content	< 1 x 10 ⁶ /unit	1%
pH measured (+22 C) at the end of the recommended shelf life	> 6.4	4 per month

General Specification:

Labelling:

See Appendix I

IBTS/PMF/SPEC/0214		Ver 7	Page 4 of 6	
Storage:	stored at 22°C	Platelets Adult Dose with Plasma / PAS Irradiated (69740) should be stored at $22^{\circ}C \pm 2^{\circ}C$. The total storage time is 7 days when stored with continuous gentle agitation.		
Irradiation:	Platelets are i	rradiated routinely during	preparation.	
Transportation:	Transport containers should be kept open at room temperature for 30 minutes before use. During transportation from the Irish Blood Transfusion Service to the place where they are intended for use the temperature of platelets must be kept as close as possible to the recommended storage temperature. On receipt, unless intended for immediate therapeutic use, they should be transferred to storage with continuous gentle agitation on a device approved for the purpose, at 22°C $\pm 2^{\circ}$ C.			
Indications for Use:	1 1	latelet replacement where s causing significant haem	5	
Precautions in Use:	• Swirling	phenomenon must be dem	onstrated before infusion.	
		must be infused intravenou g an in line 170 – 200 μm	usly through a fresh infusion set filter.	
Ner Ir			child bearing potential should atelets from Rh D positive done	
erifyw		of the temperature of stora contamination is increased	ge and preparation the risk of l.	
2	• No soluti	on should be added to the	bag or giving set.	
	-	ents should be inspected vi colour or visible clots.	sually for defects, leakage,	
	• Not record	nmended in cases of plasn	na intolerance.	
Adverse Effects Incl	ude:			
	• <u>Circulato</u>	ry overload		
		tic transfusion reaction due tible plasma in the compon		
	• Non-haer	nolytic transfusion reaction	n (mainly chills fever and	

- <u>Non-haemolytic transfusion reaction</u> (mainly chills, fever and urticaria). The risk is reduced by leucodepletion.
- <u>Anaphylaxis</u>

- <u>Pathogen transmission</u>
 - Despite careful donor selection and laboratory screening procedures, infections including Syphilis, Viral Hepatitis, HIV, HTLV 1 & 11 and other viruses and protozoa (e.g. malaria) may, in rare instances, occur.
 - vCJD transmission
 - Transmission of other pathogens that are not tested for or recognised.
 - The risk of CMV transmission is minimal as the components are leucodepleted
 - Sepsis due to bacterial contamination (reduced but not eliminated by bacterial screening)
- <u>Metabolic upset</u>
 - Citrate toxicity, especially in neonates and in patients with impaired hepatic function.
- <u>Immunological effects</u>
 - Alloimunisation to HLA and HPA antigens
 - Post Transfusion purpura (PTP), especially in parous female recipients
 - The risk of Graft vs Host Disease (GvHD) in immuno compromised recipients is eliminated by irradiation
 - Transfusion related Acute Lung injury (TRALI) by donor HLA/granulocyte antibodies

Serious Adverse Reaction

Please inform the IBTS immediately about any event relating to suspected bacterial sepsis/ transfusion associated bacterial sepsis

Serious adverse reactions should be reported to:

National Haemovigilance Office

Irish Blood Transfusion Service National Blood Centre James's Street Dublin 8

AND


Quality Assurance Manager Irish Blood Transfusion Service

AT EITHER

National Blood Centre James's Street Dublin 8

OR

Munster Regional Transfusion Centre St Finbarr's Hospital Douglas Road, Cork



N.B. Stated volume for illustration purposes only.